

OBSERVATIONS ON FOOT AND MOUTH DISEASE IN VACCINATED AND UNVACCINATED WILDLIFE IN THE UNITED ARAB EMIRATES

Tom Bailey¹, Declan O'Donovan², Joerg Kinne³, Ulrich Wernery³

¹The Dubai Falcon Hospital, P.O. Box 23919, Dubai, UAE; ²Wadi al Safa Wildlife Centre, P.O. Box 27875, Dubai, UAE; ³Central Veterinary Research Laboratory, P.O. Box 597, Dubai, UAE.

Foot-and-mouth disease (FMD) is a highly contagious, but usually nonlethal disease of ruminants, characterized by vesiculation of the oral mucosa and of the skin of the feet (Thompson et al, 2001). The virus is endemic in domestic livestock populations in the Middle East (serotype O and A) (Knowles et al 2005). There are few reports of FMD in wildlife in the region, although FMD was reported in gazelles in Israel in 2007 (Promed, 2009) and Shimshony et al (1986) described a FMD outbreak in mountain gazelles that caused severe mortality.

Our article summarises the features of two outbreaks of FMD in one unvaccinated collection and a second vaccinated collection of wild ungulates and highlights the importance of vaccination protocols to protect wildlife in a region where infectious viral diseases, including FMD, are endemic.



Fig 1. Gazelle walking on tip-toe

(~45) (*Oryx leucoryx*), mountain gazelle (~8) (*Gazella gazella*), impala (~48) (*Aepyceros melampus*) and Speke's gazelle (~36) (*Gazella spekei*) held at a collection outside Dubai. Most ungulates had been vaccinated annually against FMD since 2006. The Arabian oryx, mountain gazelle and Speke's gazelle and a group of 10 sand gazelle were kept in fenced enclosures and every adult animal in these groups had been vaccinated within the previous 3 months. The blackbuck, impala and the remaining sand gazelle population lived within the 350 hectares of the walled grounds of the collection. Although many of these animals were captured and vaccinated annually by herding them into a raceway system (O'Donovan and Bailey 2006), it was not possible to catch all of them. We estimate that 75 % of these animals were vaccinated 10-12 months previously. On 29/01/2009 a post parturient female blackbuck was found depressed and unable to stand next to a dead calf that she had delivered. The female was euthanased and submitted to CVRL. Lesions observed in the female included tongue erosions, focal ulceration of the lip and white stripes on the myocardium (Fig 6). FMD virus type O was isolated from tissues. During February, FMD was confirmed in one stillborn oryx calf, one juvenile blackbuck, and 2 sand gazelle calves. In May, FMD was confirmed in a week old mountain gazelle calf. Other than the first euthanased female blackbuck, FMD has not been confirmed in other dead adult animals at site 2 from the first case until the current time (June 2009).

Viral isolates from both outbreaks were sent to the World Reference Laboratory at the Institute for Animal Health (UK). The O virus was closely related to FMD strains from India (Ind-2001) and Iran (Ir-2001).

The source of infection was not confirmed for either outbreak. At site 1, the infection may have originated from recently imported livestock or from cow dung compost, both originating from adjacent countries where FMD is known to be endemic. At site 2, the authors were informed by local veterinarians that a sheep farm adjacent to the collection had FMD immediately prior to the outbreak in the wildlife.

These outbreaks demonstrate how FMD has the potential to wreak havoc upon a susceptible population of unvaccinated exotic ungulates managed in captivity. Few wildlife collections in the Middle East routinely vaccinate their animals against infectious diseases. An important reason is that large numbers of semi-free living exotic ungulates such as gazelles and oryx are challenging to catch and safely handle. Handling systems for exotic ungulates are becoming more commonly used (Tamer, Fauna Research, USA). Since 2006 an annual vaccination programme was initiated for all ungulates at Site 2.

The abstract of a study by Kilgalon et al (2008) assessing the immunological response of Arabian oryx to FMD vaccine is presented in

Site 1 (unvaccinated) was a collection of dorcas gazelles (~100) (*Gazella dorcas*) and sika deer (~25) (*Cervus nippon*) living within a large walled garden within Dubai. A dairy farm was located within the grounds. The week before the investigation many gazelles had died. On 29/12/2008 it was noted that the majority of deer and gazelles were lame (see Figures). The gazelle either walked on tip-toe with arched backs (Fig 1), or exhibited unilateral 'carrying leg' lameness in just one limb (Fig 2), some animals were immobile and when approached they were easily caught and examined. Many young gazelles died. Although most of the deer were observed to be lame, none died. Four immobile gazelle were euthanased and submitted to the CVRL. Approximately 50% of the gazelle died over a 6 week period. Tongue erosions were observed, but foot lesions were not observed. FMD virus type O was isolated from tongue lesions, and organs.

Site 2 was a private collection comprising blackbuck (~80) (*Antelope cervicapra*), sand gazelle (~20) (*Gazella subgutturosa marica*), Arabian oryx, mountain gazelle and Speke's gazelle and a group of 10 sand gazelle were kept in fenced enclosures and every adult animal in these groups had been vaccinated within the previous 3 months. The blackbuck, impala and the remaining sand gazelle population lived within the 350 hectares of the walled grounds of the collection. Although many of these animals were captured and vaccinated annually by herding them into a raceway system (O'Donovan and Bailey 2006), it was not possible to catch all of them. We estimate that 75 % of these animals were vaccinated 10-12 months previously. On 29/01/2009 a post parturient female blackbuck was found depressed and unable to stand next to a dead calf that she had delivered. The female was euthanased and submitted to CVRL. Lesions observed in the female included tongue erosions, focal ulceration of the lip and white stripes on the myocardium (Fig 6). FMD virus type O was isolated from tissues. During February, FMD was confirmed in one stillborn oryx calf, one juvenile blackbuck, and 2 sand gazelle calves. In May, FMD was confirmed in a week old mountain gazelle calf. Other than the first euthanased female blackbuck, FMD has not been confirmed in other dead adult animals at site 2 from the first case until the current time (June 2009).



Fig 2. Gazelle showing unilateral lameness



Fig 3. Mouth lesions in a gazelle with foot and mouth disease

this newsletter. Kilgallon et al (2008) concluded that a single dose of FMD vaccine may not elicit a sufficient antibody response in Arabian oryx to confer lasting protection and recommended as a standard prophylactic regimen, follow up doses as in domestic livestock. However, from a practical perspective, capturing exotic hoofstock for booster injections one month after primary vaccination, or even biannually is practically impossible in most collections. Our observations indicate that, although most animals at site 2 were only vaccinated annually, they were afforded good protection when exposed to the same FMD strain that caused high mortality in unvaccinated gazelle at site 1. Clearly, further research is required in this area.

References

O'Donovan, D. and Bailey, T. (2006) Restraint of Arabian oryx (*Oryx leucoryx*) in Dubai, United Arab Emirates using a mobile raceway, Proceedings of the 2nd Conference of the International Congress of Zookeepers: Gold Coast, Australia.

Kilgallon, C., Bailey, T.A., O'Donovan, D. Wernery, U. and Alexandersen, S. (2008). A Temporal Assessment of Seroconversion in Response to Inactivated Foot and Mouth Disease Vaccine in Arabian Oryx (*Oryx leucoryx*). *Veterinary Record*. 163: 717-720.

Knowles, N.J., Samuel, A.R., Davies, P.R., Midgley, R.J., and Valarcher, J.F. (2005) Pandemic Strain of Foot and Mouth Disease Virus Serotype O. *Emerging Infectious Diseases*, Vol 11 (12). Pp 1887-1893

Promed FOOT & MOUTH DISEASE - ISRAEL (02): OIE, SEROTYPE O, SPREAD. Date: Sun 15 Feb 2009 Source: OIE/WAHID Weekly Disease Information Vol. 22 - No. 08, February 2009 <http://www.oie.int/wahis/public.php?page=single_report&pop=1&reportid=7791>

Shimshony, A., Orgad, U., Baharav, D., Prudovsky, S., Yakobson, B., Bar Moshe, B. and Dagan, D. (1986) Malignant foot-and-mouth disease in mountain gazelles. *Veterinary Record* 19: 175-176.

Thompson, G.R., Bengis, R.G., Brown, C.C. 2001. Picornavirus infections. In: *Infectious diseases of wild mammals* (eds Elizabeth S. Williams and Ian K. Barker). 3rd ed, Iowa State University Press. Pp 119-124.